

Relativistic Wave Equations in the Helicity Basis

V. V. Varlamov*

Abstract

The principal series of unitary representations of the Lorentz group has been considered in the helicity basis. Decompositions of tensor products of the spinspaces are studied in the framework of projective representations of the symmetric group. Higher-spin Gel'fand-Yaglom equations are defined in the helicity basis over an arbitrary representation space. Applications of decomposable and indecomposable Gel'fand-Yaglom equations to particle physics are discussed.

1 Introduction

As known [17], a root subgroup of a semisimple Lie group O_4 (a rotation group of the 4-dimensional space) is a normal divisor of O_4 . For that reason the 6-parameter group O_4 is semisimple, and is represented by a direct product of the two 3-parameter unimodular groups. By analogy with the group O_4 , a double covering $SL(2, \mathbb{C})$ of the proper orthochronous Lorentz group $\mathfrak{G}_+$ (a rotation group of the 4-dimensional Minkowski spacetime) is semisimple, and is represented by a direct product of the two 3-parameter special unimodular groups, $SL(2, \mathbb{C}) \simeq SU(2) \otimes SU(2)$. An explicit form of this isomorphism can be obtained by means of a complexification of the group $SU(2)$, that is, $SL(2, \mathbb{C}) \simeq \text{complex}(SU(2)) \simeq SU(2) \otimes SU(2)$ [44].

*Department of Mathematics, Siberia State University of Industry, Kirova 42, Novokuznetsk 654007, Russia.

Moreover, in the works [2, 12] the Lorentz group is represented by a product $SU_R(2) \otimes SU_L(2)$, and spinors

$$\psi(p^\mu) = \begin{pmatrix} \phi_R(p^\mu) \\ \phi_L(p^\mu) \end{pmatrix}$$

are transformed within $(j, 0) \oplus (0, j)$ representation space. The components $\phi_R(p^\mu)$ and $\phi_L(p^\mu)$ correspond to different helicity states (right- and left-handed spinors).

2 Helicity basis

Let $\mathfrak{g} \rightarrow T_{\mathfrak{g}}$ be an arbitrary linear representation of group $\mathfrak{G}_+$ and let $A_i(t) = T_{a_i(t)}$ be an infinitesimal operator corresponded the rotation $a_i(t) \in \mathfrak{G}_+$. Analogously, we have $B_i(t) = T_{b_i(t)}$, where $b_i(t) \in \mathfrak{G}_+$ is a hyperbolic rotation. The operators A_i and B_i satisfy the following commutation relations:

$$\left. \begin{aligned} [A_1, A_2] &= A_3, & [A_2, A_3] &= A_1, & [A_3, A_1] &= A_2, \\ [B_1, B_2] &= -A_3, & [B_2, B_3] &= -A_1, & [B_3, B_1] &= -A_2, \\ [A_1, B_1] &= 0, & [A_2, B_2] &= 0, & [A_3, B_3] &= 0, \\ [A_1, B_2] &= B_3, & [A_1, B_3] &= -B_2, \\ [A_2, B_3] &= B_1, & [A_2, B_1] &= -B_3, \\ [A_3, B_1] &= B_2, & [A_3, B_2] &= -B_1. \end{aligned} \right\} \quad (1)$$

Denoting $I^{23} = A_1$, $I^{31} = A_2$, $I^{12} = A_3$, and $I^{01} = B_1$, $I^{02} = B_2$, $I^{03} = B_3$ we can write the relations (1) in a more compact form:

$$[I^{\mu\nu}, I^{\lambda\rho}] = \delta_{\mu\rho} I^{\lambda\nu} + \delta_{\nu\lambda} I^{\mu\rho} - \delta_{\nu\rho} I^{\mu\lambda} - \delta_{\mu\lambda} I^{\nu\rho}.$$

Let us consider the operators

$$\begin{aligned} X_k &= \frac{1}{2}(A_k + iB_k), & Y_k &= \frac{1}{2}(A_k - iB_k), \\ & & (k &= 1, 2, 3). \end{aligned} \quad (2)$$

Using the relations (1) we find that

$$\begin{aligned} [X_1, X_2] &= X_3, & [X_2, X_3] &= X_1, & [X_3, X_1] &= X_2, \\ [Y_1, Y_2] &= Y_3, & [Y_2, Y_3] &= Y_1, & [Y_3, Y_1] &= Y_2, \\ [X_k, Y_l] &= 0, & (k, l &= 1, 2, 3). \end{aligned} \quad (3)$$

Further, taking

$$\left. \begin{aligned} X_+ &= X_1 + iX_2, & X_- &= X_1 - iX_2, \\ Y_+ &= Y_1 + iY_2, & Y_- &= Y_1 - iY_2 \end{aligned} \right\} \quad (4)$$

we see that in virtue of commutativity of the relations (3) a space of an irreducible finite-dimensional representation of the group $\mathfrak{G}_+$ can be stretched on the totality of $(2l+1)(2\dot{l}+1)$ basis vectors $|l, m; \dot{l}, \dot{m}\rangle$, where $l, m, \dot{l}, \dot{m}$ are integer or half-integer numbers, $-l \leq m \leq l$, $-\dot{l} \leq \dot{m} \leq \dot{l}$. Therefore,

$$\begin{aligned} X_- |l, m; \dot{l}, \dot{m}\rangle &= \sqrt{(\dot{l} + \dot{m})(\dot{l} - \dot{m} + 1)} |l, m; \dot{l}, \dot{m} - 1\rangle \quad (\dot{m} > -\dot{l}), \\ X_+ |l, m; \dot{l}, \dot{m}\rangle &= \sqrt{(\dot{l} - \dot{m})(\dot{l} + \dot{m} + 1)} |l, m; \dot{l}, \dot{m} + 1\rangle \quad (\dot{m} < \dot{l}), \\ X_3 |l, m; \dot{l}, \dot{m}\rangle &= \dot{m} |l, m; \dot{l}, \dot{m}\rangle, \\ Y_- |l, m; \dot{l}, \dot{m}\rangle &= \sqrt{(l + m)(l - m + 1)} |l, m - 1; \dot{l}, \dot{m}\rangle \quad (m > -l), \\ Y_+ |l, m; \dot{l}, \dot{m}\rangle &= \sqrt{(l - m)(l + m + 1)} |l, m + 1; \dot{l}, \dot{m}\rangle \quad (m < l), \\ Y_3 |l, m; \dot{l}, \dot{m}\rangle &= m |l, m; \dot{l}, \dot{m}\rangle. \end{aligned} \quad (5)$$

From the relations (3) it follows that each of the sets of infinitesimal operators X and Y generates the group $SU(2)$ and these two groups commute with each other. Thus, from the relations (3) and (5) it follows that the group $\mathfrak{G}_+$, in essence, is equivalent to the group $SU(2) \otimes SU(2)$. In contrast to the Gel'fand–Naimark representation for the Lorentz group [15, 26] which does not find a broad application in physics, a representation (5) is a most useful in theoretical physics (see, for example, [1, 36, 33, 34]). This representation for the Lorentz group was firstly given by Van der Waerden in his brilliant book [51].

As showed in [43] the spinspace

$$\mathbb{S}_{2^{k+r}} \simeq \underbrace{\mathbb{S}_2 \otimes \mathbb{S}_2 \otimes \cdots \otimes \mathbb{S}_2}_{k \text{ times}} \otimes \underbrace{\dot{\mathbb{S}}_2 \otimes \dot{\mathbb{S}}_2 \otimes \cdots \otimes \dot{\mathbb{S}}_2}_{r \text{ times}} \quad (6)$$

is the minimal left ideal of the complex Clifford algebra

$$\mathbb{C}_{2n} \simeq \underbrace{\mathbb{C}_2 \otimes \mathbb{C}_2 \otimes \cdots \otimes \mathbb{C}_2}_{k \text{ times}} \otimes \underbrace{\mathbb{C}_2^* \otimes \mathbb{C}_2^* \otimes \cdots \otimes \mathbb{C}_2^*}_{r \text{ times}}. \quad (7)$$

The algebras $\mathbb{C}_2$ ($\mathbb{C}_2^*$) and the spinspace $\mathbb{S}_2$ ($\dot{\mathbb{S}}_2$) correspond to fundamental representations $\tau_{\frac{1}{2},0}$ ($\tau_{0,\frac{1}{2}}$) of the Lorentz group $\mathfrak{G}_+$.

In general case the spinspace (6) is reducible, that is, there exists a decomposition of the original spinspace $\mathbb{S}_{2^{k+r}}$ into a direct sum of irreducible subspaces with respect to a representation

$$\underbrace{\tau_{\frac{1}{2},0} \otimes \tau_{\frac{1}{2},0} \otimes \cdots \otimes \tau_{\frac{1}{2},0}}_{k \text{ times}} \otimes \underbrace{\tau_{0,\frac{1}{2}} \otimes \tau_{0,\frac{1}{2}} \otimes \cdots \otimes \tau_{0,\frac{1}{2}}}_{r \text{ times}}. \quad (8)$$

First of all, let us consider a decomposition of the spinspace

$$\mathbb{S}_{2^m} \simeq \mathbb{S}_2 \otimes \mathbb{S}_2 \otimes \cdots \otimes \mathbb{S}_2 \quad (9)$$

into irreducible subspaces with respect to an action of the representation

$$\tau_{\frac{1}{2},0} \otimes \tau_{\frac{1}{2},0} \otimes \cdots \otimes \tau_{\frac{1}{2},0}. \quad (10)$$

2.1 Projective representations of the symmetric group

A group ring $\mathbb{Z}(S_m)$ and a group algebra $\mathbb{C}(S_m)$ of the symmetric group S_m act in the spinspace (9). In 1911 [35] (see also [25, 39]), Schur showed that over the field $\mathbb{F} = \mathbb{C}$ there exists a nontrivial $\mathbb{Z}_2$ -extension T_n of the symmetric group S_n defined by the sequence

$$1 \longrightarrow \mathbb{Z}_2 \longrightarrow T_n \longrightarrow S_n \longrightarrow 1,$$

where

$$\begin{aligned} T_n = \{ & z, t_1, \dots, t_{n-1} : z^2 = 1, zt_i = t_i z, \\ & t_i^2 = z, (t_j t_{j+1})^3 = z, t_k t_l = z t_l t_k \} \\ & 1 \leq i \leq n-1, 1 \leq j \leq n-2, k \leq l-2. \end{aligned}$$

The group T_n has order $2(n!)$. The subgroup $\mathbb{Z}_2 = \{1, z\}$ is central and, is contained in the commutator subgroup of T_n , $T_n/\mathbb{Z}_2 \simeq S_n$ ($n \geq 4$). If $n < 4$, then every projective representation of S_n is projectively equivalent to a linear representation.

Spinor representations of the transpositions t_k are defined by elements of the group algebra $\mathbb{C}(S_n)$:

$$t_k = \sqrt{\frac{k-1}{2k}} \mathcal{E}_{k-1} - \sqrt{\frac{k+1}{2k}} \mathcal{E}_k, \quad k = 1, \dots, m,$$

where

$$\begin{aligned}
\mathcal{E}_1 &= \sigma_1 \otimes \mathbf{1}_2 \otimes \cdots \otimes \mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2, \\
\mathcal{E}_2 &= \sigma_3 \otimes \sigma_1 \otimes \mathbf{1}_2 \otimes \cdots \otimes \mathbf{1}_2 \otimes \mathbf{1}_2, \\
\mathcal{E}_3 &= \sigma_3 \otimes \sigma_3 \otimes \sigma_1 \otimes \mathbf{1}_2 \otimes \cdots \otimes \mathbf{1}_2, \\
&\dots\dots\dots \\
\mathcal{E}_m &= \sigma_3 \otimes \sigma_3 \otimes \cdots \otimes \sigma_3 \otimes \sigma_1, \\
\mathcal{E}_{m+1} &= \sigma_2 \otimes \mathbf{1}_2 \otimes \cdots \otimes \mathbf{1}_2, \\
\mathcal{E}_{m+2} &= \sigma_3 \otimes \sigma_2 \otimes \mathbf{1}_2 \otimes \cdots \otimes \mathbf{1}_2, \\
&\dots\dots\dots \\
\mathcal{E}_{2m} &= \sigma_3 \otimes \sigma_3 \otimes \cdots \otimes \sigma_3 \otimes \sigma_2.
\end{aligned} \tag{11}$$

are the tensor products of m Pauli matrices. These matrices form a spinor basis of the complex Clifford algebra $\mathbb{C}_n$ ($n = 2m$) in the Brauer-Weyl representation [7]. At this point there is an isomorphism $\mathbb{C}_{2m} \simeq \mathbb{M}_{2^m}(\mathbb{C})$, where $\mathbb{M}_{2^m}(\mathbb{C})$ is a full matrix algebra over the field $\mathbb{F} = \mathbb{C}$. If $n = 2m + 1$ we add one more matrix

$$\mathcal{E}_{2m+1} = \underbrace{\sigma_3 \otimes \sigma_3 \otimes \cdots \otimes \sigma_3}_{m \text{ times}}.$$

In virtue of an isomorphism $\mathbb{C}_{2m+1} \simeq \mathbb{C}_{2m} \oplus \mathbb{C}_{2m}$ the units of $\mathbb{C}_{2m+1}$ are represented by direct sums $\mathcal{E}_1 \oplus \mathcal{E}_1, \mathcal{E}_2 \oplus \mathcal{E}_2, \dots, \mathcal{E}_{2m} \oplus \mathcal{E}_{2m}$. Besides, there exists a mapping $\epsilon : \mathbb{C}_{2m+1} \rightarrow \mathbb{C}_{2m}$. In the result of the homomorphism ϵ we have a quotient algebra ${}^\epsilon\mathbb{C}_{2m} \simeq \mathbb{C}_{2m+1} / \text{Ker } \epsilon$, where $\text{Ker } \epsilon = \{\mathcal{A}^1 - \varepsilon\omega\mathcal{A}^1\}$ is a kernel of ϵ , $\mathcal{A}^1$ is an arbitrary element of the subalgebra $\mathbb{C}_{2m}$, ω is a volume element of $\mathbb{C}_{2m+1}$ [43].

Recently, projective representations of the symmetric group have been used at the study of fractional quantum Hall effect [29, 52, 53] and non-Abelian statistics [13].

In accordance with a general Weyl scheme [54] a decomposition of tensor products of representations of the groups $U(n)$ ($SU(n)$) is realized via a decomposition of a module of the group algebra $\mathbb{C}(S_m)$, $\mathbf{1} = f(\alpha_1) + f(\alpha_2) + \dots + f(\alpha_s)$, where $f(\alpha_i)$ are the primitive idempotents of $\mathbb{C}(S_m)$ (Young symmetrizers), α_i are the Young diagrams, ($i = 1, \dots, s$). The Weyl scheme fully admits the Schur's $\mathbb{Z}_2$ -extension T_n , but an explicit form of the Young's orthogonal representation has been unanswered so far [27, 28].

2.2 Decomposition of the spinspace

The decomposition of the spinspace (9) with respect to $SL(2, \mathbb{C})$ is a simplest case of the Weyl scheme. Every irreducible representation of the group $SL(2, \mathbb{C})$ is defined by the Young diagram consisting of only one row. For that reason the representation (10) is realized in the space $\text{Sym}_{(m,0)}$ of all symmetric spintensors of the rank m . Dimension of $\text{Sym}_{(m,0)}$ is equal to $m + 1$.

In its turn, every element of the spinspace (6), related with the representation (8), corresponds to an element of $\mathbb{S}_{2^k} \otimes \mathbb{S}_{2^r}$ (representations $\boldsymbol{\tau}_{\frac{k}{2},0} \otimes \boldsymbol{\tau}_{\frac{r}{2},0}$ and $\boldsymbol{\tau}_{\frac{k}{2},0} \otimes \boldsymbol{\tau}_{0,\frac{r}{2}}$ are equivalent). This equivalence can be described as follows

$$\varphi \otimes \psi \longrightarrow \varphi \otimes \psi I, \quad (12)$$

where $\varphi, \psi \in \mathbb{S}_{2^k}$, $\psi I \in \mathbb{S}_{2^r}$ and

$$I = \lambda \begin{pmatrix} 0 & & -1 \\ & \ddots & \\ (-1)^{\frac{r+k}{2}} & & 0 \end{pmatrix}$$

is the matrix of a bilinear form (this matrix is symmetric if $\frac{r+k}{2} \equiv 0 \pmod{2}$ and skewsymmetric if $\frac{r+k}{2} \equiv 1 \pmod{2}$). In such a way, the representation (8) is realized in a symmetric space $\text{Sym}_{(k,r)}$ of dimension $(k+1)(r+1)$ (or $(2l_1+1)(2l_2+1)$ if suppose $l_1 = k/2$ and $l_2 = r/2$). The decomposition of (8) is given by a Clebsh-Gordan formula

$$\boldsymbol{\tau}_{l_1 l'_1} \otimes \boldsymbol{\tau}_{l_2 l'_2} = \sum_{|l_1 - l_2| \leq k \leq l_1 + l_2; |l'_1 - l'_2| \leq k' \leq l'_1 + l'_2} \boldsymbol{\tau}_{kk'}.$$

where the each $\boldsymbol{\tau}_{kk'}$ acts in the space $\text{Sym}_{(k,k')}$. In its turn, every space $\text{Sym}_{(k,r)}$ can be represented by a space of polynomials

$$p(z_0, z_1, \bar{z}_0, \bar{z}_1) = \sum_{\substack{(\alpha_1, \dots, \alpha_k) \\ (\dot{\alpha}_1, \dots, \dot{\alpha}_r)}} \frac{1}{k! r!} a^{\alpha_1 \dots \alpha_k \dot{\alpha}_1 \dots \dot{\alpha}_r} z_{\alpha_1} \dots z_{\alpha_k} \bar{z}_{\dot{\alpha}_1} \dots \bar{z}_{\dot{\alpha}_r} \quad (13)$$

$$(\alpha_i, \dot{\alpha}_i = 0, 1)$$

with the basis (the basis of a representation $(l_1 l_2) \oplus (l_2 l_1)$)

$$\begin{aligned} z_{m_1 m_2}^{l_1 l_2} &= \frac{\zeta_1^{l_1+m_1} \zeta_2^{l_1-m_1} \zeta_1^{l_2+m_2} \zeta_2^{l_2-m_2} (\zeta_1 \zeta_2' - \zeta_2 \zeta_1')^{2l_1} (\zeta_1 \zeta_2' - \zeta_2 \zeta_1')^{2l_2}}{\sqrt{(l_1+m_1)!(l_1-m_1)!(l_2+m_2)!(l_2-m_2)!}}, \\ z_{m_2 m_1}^{l_2 l_1} &= \frac{\zeta_1^{l_1+m_1} \zeta_2^{l_1-m_1} \zeta_1^{l_2+m_2} \zeta_2^{l_2-m_2} (\zeta_1 \zeta_2' - \zeta_2 \zeta_1')^{2l_1} (\zeta_1 \zeta_2' - \zeta_2 \zeta_1')^{2l_2}}{\sqrt{(l_1+m_1)!(l_1-m_1)!(l_2+m_2)!(l_2-m_2)!}}. \end{aligned}$$

The vectors $z_{mm'}^{ll'}$ of the canonical helicity basis have the form

$$z_{mm'}^{ll'} = \sum_{j+k=m, j'+k'=m'} C(l_1, l_2, l; j, k, m) C(l_1', l_2', l'; j', k', m') \zeta_{jj'} \otimes \zeta_{kk'}, \quad (14)$$

where

$$C(l_1, l_2, l; j, k, m) C(l_1', l_2', l'; j', k', m') = B_{l_1, l_2, l; l_1', l_2', l'}^{j, k, m; j', k', m'}$$

are the Clebsch–Gordan coefficients of the group $SL(2, \mathbb{C})$. Expressing the Clebsch–Gordan coefficients $C(l_1, l_2, l; j, k, j+k)$ of the group $SU(2)$ via a generalized hypergeometric function ${}_3F_2$ (see, for example, [32, 24, 47, 50]) we see that CG-coefficients of $SL(2, \mathbb{C})$ have the form

$$\begin{aligned} B_{l_1, l_2, l; l_1', l_2', l'}^{j, k, m; j', k', m'} &= (-1)^{l_1+l_1'-j-j'} \times \\ &\quad \frac{\Gamma(l_1+l_2-m+1)\Gamma(l_1'+l_2'-m'+1)}{\Gamma(l_2-l_1+m+1)\Gamma(l_2'-l_1'+m'+1)} \times \\ &\quad \left(\frac{(l-m)!(l+l_2-l_1)!(l_1-j)!(l_2+k)!(l+m)!(2l+1)}{(l_1-l_2+l)!(l_1+l_2-l)!(l_1+l_2+l)!(l_1-j)!(l_2-k)!} \right)^{\frac{1}{2}} \times \\ &\quad \left(\frac{(l'-m')!(l'+l_2'-l_1')!(l_1'-j')!(l_2'+k')!(l'+m')!(2l'+1)}{(l_1'-l_2'+l')!(l_1'+l_2'-l')!(l_1'+l_2'+l')!(l_1'-j')!(l_2'-k')!} \right)^{\frac{1}{2}} \times \\ &\quad {}_3F_2 \left(\begin{matrix} l+m+1, -l+m, -l_1+j \\ -l_1-l_2+m, l_2-l_1+m+1 \end{matrix} \middle| 1 \right) \times \\ &\quad {}_3F_2 \left(\begin{matrix} l'+m'+1, -l'+m', -l_1'+j' \\ -l_1'-l_2'+m', l_2'-l_1'+m'+1 \end{matrix} \middle| 1 \right), \quad (15) \end{aligned}$$

where $m = j+k$, $m' = j'+k'$.

Infinitesimal operators of $\mathfrak{G}_+$ in the helicity basis have a very simple form

$$\begin{aligned}
A_1 &= -\frac{i}{2}\alpha_m^l \xi_{m-1} - \frac{i}{2}\alpha_{m+1}^l \xi_{m+1}, \\
A_2 &= \frac{1}{2}\alpha_m^l \xi_{m-1} - \frac{1}{2}\alpha_{m+1}^l \xi_{m+1}, \\
A_3 &= -im\xi_m,
\end{aligned} \tag{16}$$

$$\begin{aligned}
B_1 &= \frac{1}{2}\alpha_m^l \xi_{m-1} + \frac{1}{2}\alpha_{m+1}^l \xi_{m+1}, \\
B_2 &= \frac{i}{2}\alpha_m^l \xi_{m-1} - \frac{i}{2}\alpha_{m+1}^l \xi_{m+1}, \\
B_3 &= m\xi_m,
\end{aligned} \tag{17}$$

$$\begin{aligned}
\tilde{A}_1 &= \frac{i}{2}\alpha_m^l \xi_{m-1} + \frac{i}{2}\alpha_{m+1}^l \xi_{m+1}, \\
\tilde{A}_2 &= -\frac{1}{2}\alpha_m^l \xi_{m-1} + \frac{1}{2}\alpha_{m+1}^l \xi_{m+1}, \\
\tilde{A}_3 &= im\xi_m,
\end{aligned} \tag{18}$$

$$\begin{aligned}
\tilde{B}_1 &= -\frac{1}{2}\alpha_m^l \xi_{m-1} - \frac{1}{2}\alpha_{m+1}^l \xi_{m+1}, \\
\tilde{B}_2 &= -\frac{i}{2}\alpha_m^l \xi_{m-1} + \frac{i}{2}\alpha_{m+1}^l \xi_{m+1}, \\
\tilde{B}_3 &= -m\xi_m,
\end{aligned} \tag{19}$$

where

$$\alpha_m^l = \sqrt{(l+m)(l-m+1)}.$$

2.3 Gel'fand-Naimark basis

There exists another representation basis for the Lorentz group

$$\begin{aligned}
H_3 \xi_{lm} &= m \xi_{lm}, \\
H_+ \xi_{lm} &= \sqrt{(l+m+1)(l-m)} \xi_{l,m+1}, \\
H_- \xi_{lm} &= \sqrt{(l+m)(l-m+1)} \xi_{l,m-1}, \\
F_3 \xi_{lm} &= C_l \sqrt{l^2 - m^2} \xi_{l-1,m} - A_l m \xi_{l,m} - \\
&\quad - C_{l+1} \sqrt{(l+1)^2 - m^2} \xi_{l+1,m}, \\
F_+ \xi_{lm} &= C_l \sqrt{(l-m)(l-m-1)} \xi_{l-1,m+1} - \\
&\quad - A_l \sqrt{(l-m)(l+m+1)} \xi_{l,m+1} + \\
&\quad + C_{l+1} \sqrt{(l+m+1)(l+m+2)} \xi_{l+1,m+1}, \\
F_- \xi_{lm} &= -C_l \sqrt{(l+m)(l+m-1)} \xi_{l-1,m-1} - \\
&\quad - A_l \sqrt{(l+m)(l-m+1)} \xi_{l,m-1} - \\
&\quad - C_{l+1} \sqrt{(l-m+1)(l-m+2)} \xi_{l+1,m-1}, \\
A_l &= \frac{il_0 l_1}{l(l+1)}, \quad C_l = \frac{i}{l} \sqrt{\frac{(l^2 - l_0^2)(l^2 - l_1^2)}{4l^2 - 1}}, \\
m &= -l, -l+1, \dots, l-1, l, \\
l &= l_0, l_0+1, \dots,
\end{aligned}$$

where l_0 is positive integer or half-integer number, l_1 is an arbitrary complex number. These formulas define a finite-dimensional representation of the group $\mathfrak{G}_+$ when $l_1^2 = (l_0 + p)^2$, p is some natural number. In the case $l_1^2 \neq (l_0 + p)^2$ we have an infinite-dimensional representation of $\mathfrak{G}_+$. The operators $H_3, H_+, H_-, F_3, F_+, F_-$ are

$$\begin{aligned}
H_+ &= i\mathbf{A}_1 - \mathbf{A}_2, & H_- &= i\mathbf{A}_1 + \mathbf{A}_2, & H_3 &= i\mathbf{A}_3, \\
F_+ &= i\mathbf{B}_1 - \mathbf{B}_2, & F_- &= i\mathbf{B}_1 + \mathbf{B}_2, & F_3 &= i\mathbf{B}_3.
\end{aligned}$$

This basis was firstly given by Gel'fand in 1944 (see also [18, 14, 26]). The following relations between generators $\mathbf{Y}_\pm, \mathbf{X}_\pm, \mathbf{Y}_3, \mathbf{X}_3$ and $H_\pm, F_\pm, H_3, F_3$

define a relationship between the helicity (Van der Waerden) and Gel'fand-Naimark bases

$$\begin{aligned} Y_+ &= -\frac{1}{2}(F_+ + iH_+), & X_+ &= \frac{1}{2}(F_+ - iH_+), \\ Y_- &= -\frac{1}{2}(F_- + iH_-), & X_- &= \frac{1}{2}(F_- - iH_-), \\ Y_3 &= -\frac{1}{2}(F_3 + iH_3), & X_3 &= \frac{1}{2}(F_3 - iH_3). \end{aligned}$$

2.4 Complexification of $SU(2)$

As noted previously the explicit form of the isomorphism $SL(2, \mathbb{C}) \simeq SU(2) \otimes SU(2)$ can be obtained via the complexification of $SU(2)$. Indeed, the group $SL(2, \mathbb{C})$ is a group of all complex matrices

$$\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$$

of 2-nd order with the determinant $\alpha\delta - \gamma\beta = 1$. The group $SU(2)$ is one of the real forms of $SL(2, \mathbb{C})$. The transition from $SU(2)$ to $SL(2, \mathbb{C})$ is realized via the complexification of three real parameters φ, θ, ψ (Euler angles). Let $\theta^c = \theta - i\tau, \varphi^c = \varphi - i\epsilon, \psi^c = \psi - i\varepsilon$ be complex Euler angles, where

$$\begin{aligned} 0 &\leq \operatorname{Re} \theta^c = \theta \leq \pi, & -\infty &< \operatorname{Im} \theta^c = \tau < +\infty, \\ 0 &\leq \operatorname{Re} \varphi^c = \varphi < 2\pi, & -\infty &< \operatorname{Im} \varphi^c = \epsilon < +\infty, \\ -2\pi &\leq \operatorname{Re} \psi^c = \psi < 2\pi, & -\infty &< \operatorname{Im} \psi^c = \varepsilon < +\infty. \end{aligned}$$

Replacing in $SU(2)$ the angles φ, θ, ψ by the complex angles $\varphi^c, \theta^c, \psi^c$ we come to the following matrix

$$\begin{aligned} \mathfrak{g} &= \begin{pmatrix} \cos \frac{\theta^c}{2} e^{\frac{i(\varphi^c + \psi^c)}{2}} & i \sin \frac{\theta^c}{2} e^{\frac{i(\varphi^c - \psi^c)}{2}} \\ i \sin \frac{\theta^c}{2} e^{\frac{i(\psi^c - \varphi^c)}{2}} & \cos \frac{\theta^c}{2} e^{-\frac{i(\varphi^c + \psi^c)}{2}} \end{pmatrix} = \\ &\begin{pmatrix} \left[\cos \frac{\theta}{2} \cosh \frac{\tau}{2} + i \sin \frac{\theta}{2} \sinh \frac{\tau}{2} \right] e^{\frac{\epsilon + \varepsilon + i(\varphi + \psi)}{2}} & \left[\cos \frac{\theta}{2} \sinh \frac{\tau}{2} + i \sin \frac{\theta}{2} \cosh \frac{\tau}{2} \right] e^{\frac{\epsilon - \varepsilon + i(\varphi - \psi)}{2}} \\ \left[\cos \frac{\theta}{2} \sinh \frac{\tau}{2} + i \sin \frac{\theta}{2} \cosh \frac{\tau}{2} \right] e^{\frac{\varepsilon - \epsilon + i(\psi - \varphi)}{2}} & \left[\cos \frac{\theta}{2} \cosh \frac{\tau}{2} + i \sin \frac{\theta}{2} \sinh \frac{\tau}{2} \right] e^{\frac{-\epsilon - \varepsilon - i(\varphi + \psi)}{2}} \end{pmatrix}, \end{aligned} \tag{20}$$

since $\cos \frac{1}{2}(\theta - i\tau) = \cos \frac{\theta}{2} \cosh \frac{\tau}{2} + i \sin \frac{\theta}{2} \sinh \frac{\tau}{2}$, and $\sin \frac{1}{2}(\theta - i\tau) = \sin \frac{\theta}{2} \cosh \frac{\tau}{2} - i \cos \frac{\theta}{2} \sinh \frac{\tau}{2}$. It is easy to verify that the matrix (20) coincides with a matrix of the fundamental representation of the group $SL(2, \mathbb{C})$ (in Euler parametrization):

$$\mathfrak{g}(\varphi, \epsilon, \theta, \tau, \psi, \varepsilon) = \begin{pmatrix} e^{i\frac{\varphi}{2}} & 0 \\ 0 & e^{-i\frac{\varphi}{2}} \end{pmatrix} \begin{pmatrix} e^{\frac{\epsilon}{2}} & 0 \\ 0 & e^{-\frac{\epsilon}{2}} \end{pmatrix} \begin{pmatrix} \cos \frac{\theta}{2} & i \sin \frac{\theta}{2} \\ i \sin \frac{\theta}{2} & \cos \frac{\theta}{2} \end{pmatrix} \begin{pmatrix} \cosh \frac{\tau}{2} & \sinh \frac{\tau}{2} \\ \sinh \frac{\tau}{2} & \cosh \frac{\tau}{2} \end{pmatrix} \begin{pmatrix} e^{i\frac{\psi}{2}} & 0 \\ 0 & e^{-i\frac{\psi}{2}} \end{pmatrix} \begin{pmatrix} e^{\frac{\varepsilon}{2}} & 0 \\ 0 & e^{-\frac{\varepsilon}{2}} \end{pmatrix}.$$

Moreover, the complexification of $SU(2)$ gives us the most simple and direct way for calculation of matrix elements of the Lorentz group. It is known that these elements have a great importance in quantum field theory and widely used at the study of relativistic amplitudes. The most degenerate representation of such the elements was firstly obtained by Dolginov, Toptygin and Moskalev [9, 10, 11] via an analytic continuation of representations of the group O_4 . Later on, matrix elements are studied on the hyperboloid [48, 46], on the direct product of the hyperboloid and sphere [22, 21]. The matrix elements of the principal series are studied by Ström [40, 41] in the Gel'fand-Naimark basis. However, all the matrix elements, calculated in the GN-basis, have a very complicated form.

Matrix elements of $SL(2, \mathbb{C})$ in the helicity basis (see [44]) are

$$\begin{aligned}
\mathfrak{M}_{mn}^l(\mathfrak{g}) &= e^{-m(\epsilon+i\varphi)-n(\epsilon+i\psi)} Z_{mn}^l = e^{-m(\epsilon+i\varphi)-n(\epsilon+i\psi)} \times \\
&\sum_{k=-l}^l i^{m-k} \sqrt{\Gamma(l-m+1)\Gamma(l+m+1)\Gamma(l-k+1)\Gamma(l+k+1)} \times \\
&\cos^{2l} \frac{\theta}{2} \tan^{m-k} \frac{\theta}{2} \times \\
&\sum_{j=\max(0, k-m)}^{\min(l-m, l+k)} \frac{i^{2j} \tan^{2j} \frac{\theta}{2}}{\Gamma(j+1)\Gamma(l-m-j+1)\Gamma(l+k-j+1)\Gamma(m-k+j+1)} \times \\
&\sqrt{\Gamma(l-n+1)\Gamma(l+n+1)\Gamma(l-k+1)\Gamma(l+k+1)} \cosh^{2l} \frac{\tau}{2} \tanh^{n-k} \frac{\tau}{2} \times \\
&\sum_{s=\max(0, k-n)}^{\min(l-n, l+k)} \frac{\tanh^{2s} \frac{\tau}{2}}{\Gamma(s+1)\Gamma(l-n-s+1)\Gamma(l+k-s+1)\Gamma(n-k+s+1)}.
\end{aligned} \tag{21}$$

We will call the functions Z_{mn}^l in (21) as *hyperspherical functions*. The hyperspherical functions Z_{mn}^l can be written via the hypergeometric series:

$$\begin{aligned}
Z_{mn}^l &= \cos^{2l} \frac{\theta}{2} \cosh^{2l} \frac{\tau}{2} \sum_{k=-l}^l i^{m-k} \tan^{m-k} \frac{\theta}{2} \tanh^{n-k} \frac{\tau}{2} \times \\
&{}_2F_1 \left(\begin{matrix} m-l+1, 1-l-k \\ m-k+1 \end{matrix} \middle| i^2 \tan^2 \frac{\theta}{2} \right) {}_2F_1 \left(\begin{matrix} n-l+1, 1-l-k \\ n-k+1 \end{matrix} \middle| \tanh^2 \frac{\tau}{2} \right).
\end{aligned} \tag{22}$$

Therefore, relativistic spherical functions can be expressed by means of the function (a *generalized hyperspherical function*)

$$\mathfrak{M}_{mn}^l(\mathfrak{g}) = e^{-m(\epsilon+i\varphi)} Z_{mn}^l e^{-n(\epsilon+i\psi)}, \tag{23}$$

where

$$Z_{mn}^l = \sum_{k=-l}^l P_{mk}^l(\cos \theta) \mathfrak{P}_{kn}^l(\cosh \tau), \tag{24}$$

here $P_{mn}^l(\cos \theta)$ is a generalized spherical function on the group $SU(2)$ (see [15]), and $\mathfrak{P}_{mn}^l$ is an analog of the generalized spherical function for the group $QU(2)$ (so-called Jacobi function [49]). $QU(2)$ is a group of quasiunitary unimodular matrices of second order. As well as the group $SU(2)$, the group $QU(2)$ is one of the real forms of $SL(2, \mathbb{C})$ ($QU(2)$ is noncompact). Further, from (22) we see that the function Z_{mn}^l depends on two variables θ and τ . Therefore, using Bateman factorization we can express the hyperspherical functions Z_{mn}^l via Appell functions F_1 – F_4 (hypergeometric series of two variables [4, 6]).

3 Gel'fand-Yaglom equations in the helicity basis

As a direct consequence of the isomorphism $SL(2, \mathbb{C}) \simeq SU(2) \otimes SU(2)$ we have equations

$$\begin{aligned} \sum_{i=1}^3 \Lambda_i \frac{\partial \psi}{\partial x_i} - i \sum_{i=1}^3 \Lambda_i \frac{\partial \psi}{\partial x_i^*} + \kappa^c \psi &= 0, \\ \sum_{i=1}^3 \Lambda_i^* \frac{\partial \dot{\psi}}{\partial x_i} + i \sum_{i=1}^3 \Lambda_i^* \frac{\partial \dot{\psi}}{\partial x_i^*} + \kappa^c \dot{\psi} &= 0, \end{aligned} \quad (25)$$

with invariance conditions

$$\begin{aligned} \sum_k g_{ik}^- T_{\mathfrak{g}} \Lambda_k T_{\mathfrak{g}}^{-1} &= \Lambda_i, \\ \sum_k g_{ik}^+ T_{\mathfrak{g}}^* \Lambda_k^* T_{\mathfrak{g}}^{-1} &= \Lambda_i^*. \end{aligned} \quad (26)$$

The equations (25) act in a 3-dimensional complex space $\mathbb{C}^3$. In its turn, the space $\mathbb{C}^3$ is isometric to a 6-dimensional bivector space $\mathbb{R}^6$ (parameter space or group manifold of the Lorentz group [19, 30]).

Let us find commutation relations between the matrices Λ_i , Λ_i^* and infinitesimal operators (16), (17), (18), (19). First of all, let us present transformations $T_{\mathfrak{g}}$ ($T_{\mathfrak{g}}^*$) in the infinitesimal form, $\mathbb{I} + \mathbf{A}_i \xi + \dots$, $\mathbb{I} + \mathbf{B}_i \xi + \dots$, $\mathbb{I} + \tilde{\mathbf{A}}_i \xi + \dots$, $\mathbb{I} + \tilde{\mathbf{B}}_i \xi + \dots$. Substituting these transformations into invariance

conditions (26) we obtain with an accuracy of the terms of second order the following commutation relations

$$\begin{aligned} [\mathbf{A}_1, \Lambda_1] &= 0, & [\mathbf{A}_1, \Lambda_2] &= \Lambda_3, & [\mathbf{A}_1, \Lambda_3] &= -\Lambda_2, \\ [\mathbf{A}_2, \Lambda_1] &= -\Lambda_3, & [\mathbf{A}_2, \Lambda_2] &= 0, & [\mathbf{A}_2, \Lambda_3] &= \Lambda_1, \\ [\mathbf{A}_3, \Lambda_1] &= \Lambda_2, & [\mathbf{A}_3, \Lambda_2] &= -\Lambda_1, & [\mathbf{A}_3, \Lambda_3] &= 0. \end{aligned} \quad (27)$$

$$\begin{aligned} [\mathbf{B}_1, \Lambda_1] &= 0, & [\mathbf{B}_1, \Lambda_2] &= i\Lambda_3, & [\mathbf{B}_1, \Lambda_3] &= -i\Lambda_2, \\ [\mathbf{B}_2, \Lambda_1] &= -i\Lambda_3, & [\mathbf{B}_2, \Lambda_2] &= 0, & [\mathbf{B}_2, \Lambda_3] &= i\Lambda_1, \\ [\mathbf{B}_3, \Lambda_1] &= i\Lambda_2, & [\mathbf{B}_3, \Lambda_2] &= -i\Lambda_1, & [\mathbf{B}_3, \Lambda_3] &= 0. \end{aligned} \quad (28)$$

$$\begin{aligned} [\tilde{\mathbf{A}}_1, \Lambda_1^*] &= 0, & [\tilde{\mathbf{A}}_1, \Lambda_2^*] &= -\Lambda_3^*, & [\tilde{\mathbf{A}}_1, \Lambda_3^*] &= \Lambda_2^*, \\ [\tilde{\mathbf{A}}_2, \Lambda_1^*] &= \Lambda_3^*, & [\tilde{\mathbf{A}}_2, \Lambda_2^*] &= 0, & [\tilde{\mathbf{A}}_2, \Lambda_3^*] &= -\Lambda_1^*, \\ [\tilde{\mathbf{A}}_3, \Lambda_1^*] &= -\Lambda_2^*, & [\tilde{\mathbf{A}}_3, \Lambda_2^*] &= \Lambda_1^*, & [\tilde{\mathbf{A}}_3, \Lambda_3^*] &= 0. \end{aligned} \quad (29)$$

$$\begin{aligned} [\tilde{\mathbf{B}}_1, \Lambda_1^*] &= 0, & [\tilde{\mathbf{B}}_1, \Lambda_2^*] &= +i\Lambda_3^*, & [\tilde{\mathbf{B}}_1, \Lambda_3^*] &= -i\Lambda_2^*, \\ [\tilde{\mathbf{B}}_2, \Lambda_1^*] &= -i\Lambda_3^*, & [\tilde{\mathbf{B}}_2, \Lambda_2^*] &= 0, & [\tilde{\mathbf{B}}_2, \Lambda_3^*] &= +i\Lambda_1^*, \\ [\tilde{\mathbf{B}}_3, \Lambda_1^*] &= +i\Lambda_2^*, & [\tilde{\mathbf{B}}_3, \Lambda_2^*] &= -i\Lambda_1^*, & [\tilde{\mathbf{B}}_3, \Lambda_3^*] &= 0. \end{aligned} \quad (30)$$

Further, using the latter relations and taking into account (2) it is easy to establish commutation relations between Λ_3 and generators $\mathbf{Y}_\pm$, $\mathbf{Y}_3$, $\mathbf{X}_\pm$, $\mathbf{X}_3$:

$$\left. \begin{aligned} [\mathbf{Y}_+, [\Lambda_3, \mathbf{Y}_-]] &= 2\Lambda_3, \\ [\Lambda_3, \mathbf{Y}_3] &= 0, \\ [\Lambda_3, \mathbf{X}_-] &= 0, \\ [\Lambda_3, \mathbf{X}_+] &= 0, \\ [\Lambda_3, \mathbf{X}_3] &= 0, \end{aligned} \right\} \quad (31)$$

$$\left. \begin{aligned} [\mathbf{X}_+, [\Lambda_3^*, \mathbf{X}_-]] &= 2\Lambda_3^*, \\ [\Lambda_3^*, \mathbf{X}_3] &= 0, \\ [\Lambda_3^*, \mathbf{Y}_-] &= 0, \\ [\Lambda_3^*, \mathbf{Y}_+] &= 0, \\ [\Lambda_3^*, \mathbf{Y}_3] &= 0. \end{aligned} \right\} \quad (32)$$

From (31) and (32) it immediately follows that elements of the matrices Λ_3 and Λ_3^* are

$$\Lambda_3 : \quad \left\{ \begin{aligned} c_{l-1,l,m}^{k'k} &= c_{l-1,l}^{k'k} \sqrt{l^2 - m^2}, \\ c_{l,l,m}^{k'k} &= c_{ll}^{k'k} m, \\ c_{l+1,l,m}^{k'k} &= c_{l+1,l}^{k'k} \sqrt{(l+1)^2 - m^2}. \end{aligned} \right. \quad (33)$$

$$\Lambda_3^* : \quad \left\{ \begin{aligned} f_{i-1,i,\dot{m}}^{k'k} &= c_{i-1,i}^{k'k} \sqrt{i^2 - \dot{m}^2}, \\ f_{ii,\dot{m}}^{k'k} &= c_{ii}^{k'k} \dot{m}, \\ f_{i+1,i,\dot{m}}^{k'k} &= c_{i+1,i}^{k'k} \sqrt{(i+1)^2 - \dot{m}^2}. \end{aligned} \right. \quad (34)$$

All other elements are equal to zero. If we know the elements of Λ_3 and Λ_3^* , then the elements of Λ_1 , Λ_2 and Λ_1^* , Λ_2^* can be obtained via the relations (27)–(30).

With a view to separate the variables in (25) let us assume

$$\psi_{lm}^k = \mathbf{f}_{lmk}^{l_0}(r) \mathfrak{M}_{mn}^{l_0}(\varphi, \epsilon, \theta, \tau, 0, 0), \quad (35)$$

where $l_0 \geq l$, and $-l_0 \leq m, n \leq l_0$. The full separation of variables in (25) has been given in the recent paper [45], where the hyperspherical functions $\mathfrak{M}_{m,n}^l$ are defined on a 2-dimensional complex sphere (about the 2-dimensional complex sphere see [37]). In the result of the separation we come to a system of ordinary differential equations which depends on the

radial functions,

$$\begin{aligned}
& \sum_{k'} c_{l,l-1}^{kk'} \left[2\sqrt{l^2 - m^2} \frac{d\mathbf{f}_{l-1,m,k'}^{l_0}(r)}{dr} + \right. \\
& \quad - \frac{1}{r}(l+1)\sqrt{l^2 - m^2} \mathbf{f}_{l-1,m,k'}^{l_0}(r) + \\
& \quad + \frac{i}{r}\sqrt{(l+m)(l+m-1)}\sqrt{(l_0 + m)(l_0 - m + 1)} \mathbf{f}_{l-1,m-1,k'}^{l_0}(r) + \\
& \quad \left. + \frac{i}{r}\sqrt{(l-m)(l-m-1)}\sqrt{(l_0 + m + 1)(l_0 - m)} \mathbf{f}_{l-1,m+1,k'}^{l_0}(r) \right] + \\
& \sum_{k'} c_{ll}^{kk'} \left[2m \frac{d\mathbf{f}_{l,m,k'}^{l_0}(r)}{dr} - \frac{1}{r}m \mathbf{f}_{l,m,k'}^{l_0}(r) - \right. \\
& \quad - \frac{i}{r}\sqrt{(l+m)(l-m+1)}\sqrt{(l_0 + m)(l_0 - m + 1)} \mathbf{f}_{l,m-1,k'}^{l_0}(r) + \\
& \quad \left. + \frac{i}{r}\sqrt{(l+m+1)(l-m)}\sqrt{(l_0 + m + 1)(l_0 - m)} \mathbf{f}_{l,m+1,k'}^{l_0}(r) \right] + \\
& \sum_{k'} c_{l,l+1}^{kk'} \left[2\sqrt{(l+1)^2 - m^2} \frac{d\mathbf{f}_{l+1,m,k'}^{l_0}(r)}{dr} - \right. \\
& \quad + \frac{1}{r}l\sqrt{(l+1)^2 - m^2} \mathbf{f}_{l+1,m,k'}^{l_0}(r) - \\
& \quad - \frac{i}{r}\sqrt{(l-m+1)(l-m+2)}\sqrt{(l_0 + m)(l_0 - m + 1)} \mathbf{f}_{l+1,m-1,k'}^{l_0}(r) - \\
& \quad \left. - \frac{i}{r}\sqrt{(l+m+1)(l+m+2)}\sqrt{(l_0 + m + 1)(l_0 - m)} \mathbf{f}_{l+1,m+1,k'}^{l_0}(r) \right] + \\
& \quad + \kappa^c \mathbf{f}_{lmk}^{l_0}(r) = 0,
\end{aligned}$$

$$\begin{aligned}
& \sum_{k'} c_{l,l-1}^{kk'} \left[2\sqrt{l^2 - m^2} \frac{d\mathbf{f}_{l-1,\dot{m},k'}^{i_0}(r)}{dr^*} - \right. \\
& \quad \left. - \frac{1}{r^*} (l+1) \sqrt{l^2 - m^2} \mathbf{f}_{l-1,\dot{m},k'}^{i_0}(r) + \right. \\
& \quad \left. + \frac{i}{r^*} \sqrt{(l+\dot{m})(l+\dot{m}-1)} \sqrt{(l_0+m)(l_0-m+1)} \mathbf{f}_{l-1,\dot{m}-1,k'}^{i_0}(r) + \right. \\
& \quad \left. + \frac{i}{r^*} \sqrt{(l-\dot{m})(l-\dot{m}-1)} \sqrt{(l_0+m+1)(l_0-m)} \mathbf{f}_{l-1,\dot{m}+1,k'}^{i_0}(r) \right] + \\
& \sum_{k'} c_{l,l}^{kk'} \left[2\dot{m} \frac{d\mathbf{f}_{l,\dot{m},k'}^{i_0}(r)}{dr^*} - \frac{1}{r^*} \dot{m} \mathbf{f}_{l,\dot{m},k'}^{i_0}(r) - \right. \\
& \quad \left. - \frac{i}{r^*} \sqrt{(l+\dot{m})(l-\dot{m}-1)} \sqrt{(l_0+m)(l_0-m+1)} \mathbf{f}_{l,\dot{m}-1,k'}^{i_0}(r) + \right. \\
& \quad \left. + \frac{i}{r^*} \sqrt{(l+\dot{m}+1)(l-\dot{m})} \sqrt{(l_0+m+1)(l_0-m)} \mathbf{f}_{l,\dot{m}+1,k'}^{i_0}(r) \right] + \\
& \sum_{k'} c_{l,l+1}^{kk'} \left[2\sqrt{(l+1)^2 - m^2} \frac{d\mathbf{f}_{l+1,\dot{m},k'}^{i_0}(r)}{dr^*} + \right. \\
& \quad \left. + \frac{1}{r^*} l \sqrt{(l+1)^2 - m^2} \mathbf{f}_{l+1,\dot{m},k'}^{i_0}(r) - \right. \\
& \quad \left. - \frac{i}{r^*} \sqrt{(l-\dot{m}+1)(l-\dot{m}+2)} \sqrt{(l_0+m)(l_0-m+1)} \mathbf{f}_{l+1,\dot{m}-1,k'}^{i_0}(r) - \right. \\
& \quad \left. - \frac{i}{r^*} \sqrt{(l+\dot{m}+1)(l+\dot{m}+2)} \sqrt{(l_0+m+1)(l_0-m)} \mathbf{f}_{l+1,\dot{m}+1,k'}^{i_0}(r) \right] + \\
& \quad + \kappa^c \mathbf{f}_{l\dot{m}k'}^{i_0}(r) = 0. \quad (36)
\end{aligned}$$

This system is solvable for any spin in a class of the Bessel functions. Therefore, according to (35) relativistic wave functions are expressed via the products of cylindrical and hyperspherical functions. For that reason the wave function hardly depends on the parameters of the Lorentz group. Moreover, it allows to consider solutions of RWE as the functions on the Lorentz group. On the other hand, a group theoretical description of RWE allows

to present all the physical fields on an equal footing. Namely, all these fields are the functions on the Lorentz (Poincaré) group. Such a description corresponds to a quantum field theory on the Poincaré group introduced by Lurçat [23] (see also [5, 20, 42, 16] and references therein).

3.1 Decomposable and indecomposable RWE

In accordance with equivalence (12) and decompositions of (6) and (8) the matrix Λ_3 can be written in the form

$$\Lambda_3 = \begin{pmatrix} C^1 & & & \\ & C^2 & & 0 \\ & & \ddots & \\ 0 & & & C^s & \\ & & & & \ddots \end{pmatrix}. \quad (37)$$

The matrix Λ_3^* has the same form. C^s is a spin block. The each spin block C^s is realized in the space Sym_s . The matrix elements of the corresponded representations are expressed via the hyperspherical functions. If the spin block C^s has non-null roots, then the particle possesses the spin s [14, 3, 31]. The spin block C^s in (37) consists of the elements $c_{\tau\tau'}^s$, where τ_{l_1, l_2} and $\tau_{l'_1, l'_2}$ are interlocking irreducible representations of the Lorentz group, that is, such representations, for which $l'_1 = l_1 \pm \frac{1}{2}$, $l'_2 = l_2 \pm \frac{1}{2}$. At this point the block C^s contains only the elements $c_{\tau\tau'}^s$ corresponding to such interlocking representations τ_{l_1, l_2} , $\tau_{l'_1, l'_2}$ which satisfy the conditions

$$|l_1 - l_2| \leq s \leq l_1 + l_2, \quad |l'_1 - l'_2| \leq s \leq l'_1 + l'_2.$$

According to a de Broglie theory of fusion [8, 38] interlocking representations give rise to indecomposable RWE. Otherwise, we have decomposable equations. As known, the indecomposable RWE correspond to composite particles. A wide variety of such representations and RWE it seems to be sufficient for description of all particle world.

References

- [1] A.I. Akhiezer, V.B. Berestetskii, Quantum Electrodynamics (John Wiley & Sons, New York, 1965).

- [2] D.V. Ahluwalia, D.J. Ernst, $(j, 0) \oplus (0, j)$ covariant spinors and causal propagators based on Weinberg formalism, *Int. J. Mod. Phys.* **E2**, 397–422 (1993).
- [3] V. Amar, U. Dozzio, *Gel'fand-Yaglom Equations with Charge or Energy Density of Definite Sign*, *Nuovo Cimento* **A11**, 87–99 (1972).
- [4] P. Appell, M.J. Kampé de Fériet, *Fonctions hypéométriques et hypersphériques. Polynômes d'Hermite* (Gauthier–Villars, 1926).
- [5] H. Bacry, A. Kihlberg, *Wavefunctions on homogeneous spaces*, *J. Math. Phys.* **10**, 2132–2141 (1969).
- [6] H. Bateman, A. Erdélyi, *Higher Transcendental Functions*, vol. I (Mc Grow-Hill Book Company, New York, 1953).
- [7] R. Brauer, H. Weyl, *Spinors in n dimensions*, *Amer. J. Math.* **57**, 425–449 (1935).
- [8] L. de Broglie, *Theorie de particules a spin (methode de fusion)* (Paris, 1943).
- [9] A.Z. Dolginov, *Relativistic spherical functions*, *Zh. Ehksp. Teor. Fiz.* **30**, 746–755 (1956).
- [10] A.Z. Dolginov, I.N. Toptygin, *Relativistic spherical functions. II*, *Zh. Ehksp. Teor. Fiz.* **37**, 1441–1451 (1959).
- [11] A.Z. Dolginov, A.N. Moskalev, *Relativistic spherical functions. III*, *Zh. Ehksp. Teor. Fiz.* **37**, 1697–1707 (1959).
- [12] V.V. Dvoeglazov, *Extra Dirac equations*, *Nuovo Cimento*, **B111**, 483–496 (1996).
- [13] D. Finkelstein, A. Galiautdinov, *Cliffordons*, preprint hep-th/0005039 (2000).
- [14] I.M. Gel'fand, A.M. Yaglom, *General relativistic-invariant equations and infinite-dimensional representations of the Lorentz group*, *Zh. Ehksp. Teor. Fiz.* **18**, 703–733 (1948).
- [15] I.M. Gel'fand, R.A. Minlos, Z.Ya. Shapiro, *Representations of the Rotation and Lorentz Groups and their Applications* (Pergamon Press, Oxford, 1963).

- [16] D.M. Gitman, A.L. Shelepin, *Fields on the Poincaré Group: Arbitrary Spin Description and Relativistic Wave Equations*, Int. J. Theor. Phys. **40**(3), 603–684 (2001).
- [17] M. Goto, F. Grosshans, *Semisimple Lie Algebras* (Marcel Dekker, New York, 1978).
- [18] Harish Chandra, *Infinite irreducible representations of the Lorentz group*, Proc. Royal. Soc. London **A189**, 372–401 (1947).
- [19] V.F. Kagan, *Ueber einige Zahlensysteme, zu denen die Lorentztransformation führt*, Publ. House of Institute of Mathematics, Moscow (1926).
- [20] A. Kihlberg, *Fields on a homogeneous space of the Poincaré group*, Ann. Inst. Henri Poincaré **13**, 57–76 (1970).
- [21] G.I. Kuznetsov, M.A. Liberman, A.A. Makarov, Ya.A. Smorodinsky, *Helicity and unitary representations of the Lorentz group*, Yad. Fiz. **10**, 644–656 (1969).
- [22] M.A. Liberman, Ya.A. Smorodinsky, M.B. Sheftel, *Unitary representations of the Lorentz group and functions with spin*, Yad. Fiz. **7**, 202–214 (1968).
- [23] F. Lurçat, *Quantum field theory and the dynamical role of spin*, Physics **1**, 95 (1964).
- [24] G.Ya. Lyubarskii, *The application of group theory in physics* (Pergamon Press, Oxford, 1960).
- [25] A.O. Morris, *The spin representations of the symmetric group*, Proc. London. Math. Soc. **12** (3), 55–76 (1962).
- [26] M.A. Naimark, *Linear Representations of the Lorentz Group* (Pergamon, London, 1964).
- [27] M.L. Nazarov, *Young’s orthogonal form of irreducible projective representations of the symmetric group*, J. London Math. Soc. **42** (2), 437–451 (1990).
- [28] M.L. Nazarov, *Young’s symmetrizers for projective representations of the symmetric group*, RIMS 900-Koyoto Univ. Japan (1992).
- [29] C. Nayak, F. Wilczek, *Quantum Numbers of Hall Effect Skyrmions*, Nucl. Phys. **B479**, 529 (1996).

- [30] A.Z. Petrov, *Einstein Spaces* (Pergamon Press, Oxford, 1969).
- [31] V.A. Pletyukhov, V.I. Strazhev, *On Dirac-like relativistic wave equations*, Russian J. Phys. n.12, 38-41 (1983).
- [32] M.E. Rose, *Multipole Fields* (John Wiley & Sons, New York, 1955).
- [33] Yu.B. Rumer, A.I. Fet, *Group Theory and Quantized Fields* (Nauka, Moscow, 1977) [in Russian].
- [34] L. Ryder, *Quantum Field Theory* (Cambridge University Press, Cambridge, 1985).
- [35] I. Schur, *Über die Darstellung der symmetrischen und der alternierenden Gruppen durch gebrochene lineare substitutionen*, J. reine und angew. Math. **139**, 155–250 (1911); English translation in Int. J. Theor. Phys. **40**, 413 (2001).
- [36] S.S. Schweber, *An Introduction to Relativistic Quantum Field Theory* (Harper & Row, New York, 1961).
- [37] Ya.A. Smorodinsky, M. Huszar, *Representations of the Lorentz group and the generalization of helicity states*, Teor. Mat. Fiz. **4**, 3, 328–340 (1970).
- [38] G.A. Sokolik, *Groups Methods in the Theory of Elementary Particles* (Atomizdat, Moscow, 1965) [in Russian].
- [39] J. Stembridge, *Shifted tableaux and the projective representation of symmetric groups*, Advances in Mathematics **74**, 87–134 (1989).
- [40] S. Ström, *On the matrix elements of a unitary representation of the homogeneous Lorentz group*, Arkiv för Fysik **29**, 467–483 (1965).
- [41] S. Ström, *A note on the matrix elements of a unitary representation of the homogeneous Lorentz group*, Arkiv för Fysik **33**, 465–469 (1967).
- [42] M. Toller, *Free quantum fields on the Poincaré group*, J. Math. Phys. **37**, 2694–2730 (1996).
- [43] V.V. Varlamov, *Group Theoretical Description of Space Inversion, Time Reversal and Charge Conjugation*, preprint math-ph/0203059 (2002).

- [44] V.V. Varlamov, *Hyperspherical Functions and Linear Representations of the Lorentz Group*, preprint math-ph/0205013 (2002); to appear in Hadronic Journal.
- [45] V.V. Varlamov, *General Solutions of Relativistic Wave Equations*, preprint math-ph/0209036 (2002).
- [46] I.A. Verdiev, L.A. Dadashev, *Matrix elements of the Lorentz group unitary representation*, Yad. Fiz. **6**, 1094–1099 (1967).
- [47] I.A. Verdiev, G.A. Kerimov, Ya.A. Smorodinsky, *Clebsh-Gordan coefficients of the Lorentz group*, Yad. Fiz. **21**, 1351–1361 (1975).
- [48] N.Ya. Vilenkin, Ya.A. Smorodinsky, *Invariant expansions of relativistic amplitudes*, Zh. Ehksp. Teor. Fiz. **46**, 1793–1808 (1964).
- [49] N.Ya. Vilenkin, *Special Functions and the Theory of Group Representations* (AMS, Providence, 1968).
- [50] N.Ya. Vilenkin, A.U. Klimyk, *Representations of Lie Groups and Special Functions*, vols. 1–3. (Dordrecht: Kluwer Acad. Publ., 1991–1993).
- [51] B.L. van der Waerden, *Die Gruppentheoretische Methode in der Quantenmechanik* (Springer, Berlin, 1932).
- [52] F. Wilczek, *Some examples in the realization of symmetry*, Nucl. Phys. Proc. Suppl. **68**, 367 (1998).
- [53] F. Wilczek, *Projective Statistics and Spinors in Hilbert Space*, preprint hep-th/9806228 (1998).
- [54] H. Weyl, *Classical Groups* (Princeton, 1939).